

Catalogue of American Amphibians and Reptiles.

Russell, A.P. and A.M. Bauer. 1991. *Anolis opalinus*.

***Anolis opalinus* Gosse**
Opalescent Anole

Anolis opalinus Gosse, 1850:345. Type-locality, "Jamaica", restricted to Bluefields, Westmoreland, Jamaica by Underwood and Williams (1959). Holotype not designated (subsequently given as British Museum (Natural History) (BMNH) 1946.8.5.50, formerly 47.12.27.42), collected by P. H. Gosse, date of collection unknown (examined by authors). See Nomenclatural History.

Anolis grabami: O'Shaughnessy, 1875:273 (part).

Anolis flabellatus Cope, 1894 (1895):439. Type-locality, "Port Morant and Port Luca [Jamaica]". Holotype not designated, collected by J. Percy Moore in 1890 or 1891.

Anolis lineatopus: Barbour, 1922:668 (part).

Norops opalinus: Schwartz and Henderson, 1988:158.

- **Content.** No subspecies are currently recognized.

- **Definition.** *Anolis opalinus* is a member of the *A. grabami* series (Williams, 1976). This gracile anole is the smallest Jamaican member of the genus. Males may reach 53 mm and females 44 mm SVL, respectively (Schwartz and Henderson, 1991). Tail lengths are approximately 1.7 and 1.5 times SVL for males and females, respectively. The dorsal scales are small and granular. An erectile nuchal fold and a dorsal fold are present. The ventral scales are larger than the dorsals, smooth and slightly imbricate. The body color is typically olive to grey, with irregular mottling on the dorsum. A pale stripe runs from the labial region to the flank. The sacrum often has paired pale markings. Blue pigment may be present at the tail base. The venter is light and opalescent. Females and young exhibit dark striping on the chin and throat. Males have a burnt orange dewlap with a yellowish center and a cream to yellow border. The dewlap is rudimentary in females.

- **Descriptions.** In addition to the type descriptions of Gosse (1850) and Cope (1894 [1895]), detailed descriptions are also provided by Grant (1940), Underwood and Williams (1959), and Schwartz and Henderson (1991).

- **Illustrations.** A color illustration of the head and forebody of *A. opalinus* is included in Schwartz and Henderson (1985). Cope (1894 [1895]) depicted details of scalation. Jenssen (1978) graphed head bobbing display patterns.

- **Distribution.** *Anolis opalinus* is endemic to Jamaica. It occurs throughout the island, from sea level to over 1500 m. Barbour (1914, 1937) indicated that he considered it to be generally rare and common only around villages, due to mongoose predation else-





Figure. Male *Anolis opalinus* from Marlborough airstrip, Mandeville, Jamaica. Photograph by Anthony P. Russell.

where. It appears, however, that it may be very abundant locally (Grant, 1940; A. P. Russell, pers. obs.) and prefers mesic habitats. It is essentially arboreal but may be saxicolous where suitable habitat occurs, often occupying the crevices of dry stone walls. Individuals may be found in close proximity to human habitations, but they are not true commensals. Locality records are sparse in south central and northwestern parts of Jamaica.

- **Fossil Record.** None.

- **Pertinent Literature.** The most comprehensive systematic treatment is that of Underwood and Williams (1959), who with Grant (1940) summarized the general biology of *A. opalinus*. Hedges and Burnell (1990) used electrophoresis to examine phylogeny and biogeography. Important contributions concerning the ecology of *A. opalinus* have been made by Barbour (1914), Lazell (1966), Rand (1967a), Schoener (1970a, b), Schoener and Schoener (1971), Williams (1972, 1983), Jenssen (1973), Lister (1976a, b), and Williams and Rand (1977). Haefner (1988) analyzed the assembly rules of Jamaican anole communities, including data pertinent to *A. opalinus*. Losos (1990a, b) studied the coevolution of ecological, morphological, and behavioral attributes of members of West Indian *Anolis* communities, including *A. opalinus*. Diet and growth were studied by Floyd and Jenssen (1983, 1984) and Jenssen and Andrews (1984). Rand (1967b) and Jenssen (1978, 1979a, b) examined behavior, and Andrews and Rand (1974) discussed reproductive energetics. Immunology has



Map. Open circles indicate locality records within the range (see Distribution). The solid circle marks the restricted type-locality.

been discussed by Shochat and Dessauer (1981), and karyology by Gorman (1973) and Gorman and Atkins (1968). Fite and Lister (1981) studied vision. Telford (1975) reported on blood parasites. Lefcourt and Blaustein (1991) used data on intestinal parasite infections from Bundy et al. (1987) to test the Hamilton and Zuk hypothesis. The species was employed by Janzen and Paukstis (1991) in a study of environmental sex determination in reptiles.

• **Nomenclatural History.** Although the original description of this species clearly refers to *A. opalinus*, the putative type is not of this taxon but appears to be a juvenile specimen of *Anolis grabami* (Grant, 1940; Underwood and Williams, 1959). In a thorough discussion of the matter, Underwood and Williams (1959) pointed out that Gosse's (1850) description apparently was of a living animal and, although he referred therein to several specimens in the British Museum, never explicitly designated them as types. Much of the Jamaican material originally sent by Gosse to the British Museum had become hopelessly intermingled over the years, because even type-specimens were not labeled. *Anolis opalinus* has been confused with *A. grabami* (O'Shaughnessy, 1875; Boulenger, 1885; Barbour, 1910, 1937) and *A. lineatopus* (Barbour, 1922). These errors may stem from the incongruity between the original description and the putative type-specimen.

• **Etymology.** The name *opalinus* (L., "like an opal"), refers to the distinctive ventral coloration of the species.

Literature Cited

- Andrews, R. and A.S. Rand. 1974. Reproductive effort in anoline lizards. *Ecology* 55:1317-1327.
- Barbour, T. 1910. Notes on the herpetology of Jamaica. *Bull. Mus. Comp. Zool.* 52:273-301.
- . 1914. A contribution to the zoogeography of the West Indies, with especial reference to amphibians and reptiles. *Mem. Mus. Comp. Zool.* 44:209-359.
- . 1922. An annotated list of the Amphibia and Reptilia of Jamaica. *The Handbook of Jamaica for 1922*:666-669.
- . 1937. Third list of Antillean reptiles and amphibians. *Bull. Mus. Comp. Zool.* 82:77-166.
- Boulenger, G.A. 1885. Catalogue of the lizards in the British Museum (Natural History). 2nd ed., Vol. II. Taylor and Francis, London.
- Bundy, D.A.P., P. Vogel, and E.A. Harris. 1987. Helminth parasites of Jamaican anoles (Reptilia: Iguanidae): a comparison of the helminth fauna of 6 *Anolis* species. *J. Helminthol.* 61:77-83.
- Cope, E.D. 1894 (1895). The Batrachia and Reptilia of the University of Pennsylvania West Indian expedition of 1890 and 1891. *Proc. Acad. Nat. Sci. Philadelphia* 1894:429-442.
- Fite, K.V. and B.C. Lister. 1981. Bifoveal vision in *Anolis* lizards. *Brain Behav. Evol.* 19:144-154.
- Floyd, H.B. and T.A. Janssen. 1983. Food habits of the Jamaican lizard *Anolis opalinus*: resource partitioning and seasonal effects examined. *Copeia* 1983:319-331.
- and —. 1984. Prey diversity comparisons between stomach and hindgut of the lizard, *Anolis opalinus*. *J. Herpetol.* 18:204-205.
- Gorman, G.C. 1973. The chromosomes of the Reptilia, a cytotoxic interpretation, p. 347-424. *In* A.B. Chiarelli and E. Capanza (eds.), *Cytotaxonomy and vertebrate evolution*. Academic Press, New York.
- and L. Atkins. 1968. New karyotypic data for 16 species of *Anolis* (Sauria: Iguanidae) from Cuba, Jamaica, and the Cayman Islands. *Herpetologica* 24:13-20.
- Gosse, P.H. 1850. Descriptions of a new genus and six new species of saurian reptiles. *Ann. Mag. Nat. Hist., Ser. 2*, 6:344-348.
- Grant, C. 1940. The herpetology of Jamaica. II. The reptiles. *Bull. Inst. Jamaica, Sci. Ser.*, (1):61-148.
- Haefner, J.W. 1988. Assembly rules for Greater Antillean *Anolis* lizards: competition and random models compared. *Oecologia* 74:551-565.
- Hedges, S.B. and K.L. Burnell. 1990. The Jamaican radiation of *Anolis* (Sauria: Iguanidae): an analysis of relationships and biogeography using sequential electrophoresis. *Carib. J. Sci.* 26:31-44.
- Janzen, F.J. and G.L. Paukstis. 1991. A preliminary test of the adaptive significance of environmental sex determination in reptiles. *Evolution* 45:435-440.
- Janssen, T.A. 1973. Shift in the structural habitat of *Anolis opalinus* due to congeneric competition. *Ecology* 54:863-869.
- . 1978. Display diversity in anoline lizards and problems of interpretation, p. 269-285. *In* N. Greenberg and P.D. MacLean (eds.), *Behavior and neurology of lizards*. DHEW Publ. No. (ADM) 77-491, National Institute of Mental Health, Rockville, Maryland.
- . 1979a. Display behaviour of male *Anolis opalinus* (Sauria, Iguanidae): a case of weak display stereotypy. *Anim. Behav.* 27:173-184.
- . 1979b. Display modifiers of *Anolis opalinus* (Lacertilia: Iguanidae). *Herpetologica* 35:21-30.
- and R.M. Andrews. 1984. Seasonal growth rates in the Jamaican lizard, *Anolis opalinus*. *J. Herpetol.* 18:338-341.
- Lazell, J.D., Jr. 1966. Contributions to the herpetology of Jamaica. Studies on *Anolis reconditus* Underwood and Williams. *Bull. Inst. Jamaica, Sci. Ser.*, 18:5-15.
- Lefcourt, H. and A.R. Blaustein. 1991. Parasite load and brightness in lizards: an interspecific test of the Hamilton and Zuk hypothesis. *J. Zool. (London)* 224:491-499.
- Lister, B.C. 1976a. The nature of niche expansion in West Indian *Anolis* lizards I: ecological consequences of reduced competition. *Evolution* 30:659-676.
- . 1976b. The nature of niche expansion in West Indian *Anolis* lizards II: evolutionary components. *Evolution* 30:677-692.
- Losos, J.B. 1990a. Ecomorphology, performance capability, and scaling of West Indian *Anolis* lizards: an evolutionary analysis. *Ecol. Monogr.* 60:369-388.
- . 1990b. The evolution of form and function: morphology and locomotor performance in West Indian *Anolis* lizards. *Evolution* 44:1189-1203.
- O'Shaughnessy, A.W.E. 1875. List and revision of the species of Anolidae in the British Museum collection, with descriptions of new species. *Ann. Mag. Nat. Hist., Ser. 4*, 15:270-281.
- Rand, A.S. 1967a. The ecological distribution of the anoline lizards around Kingston, Jamaica. *Breviora* (272):1-18.
- . 1967b. Ecology and social organization in the iguanid lizard *Anolis lineatopus*. *Proc. U. S. Natl. Mus.* 122:1-79.
- Schoener, T.W. 1970a. Size patterns in West Indian *Anolis* lizards. II. Correlations with the sizes of particular sympatric species - displacement and convergence. *Amer. Nat.* 104:155-174.
- . 1970b. Nonsynchronous spatial overlap of lizards in patchy habitats. *Ecology* 51:408-418.
- and A. Schoener. 1971. Structural habitats of West Indian *Anolis* lizards I. Lowland Jamaica. *Breviora* (368):1-53.
- Schwartz, A. and R.W. Henderson. 1985. A guide to the identification of the amphibians and reptiles of the West Indies exclusive of Hispaniola. *Milwaukee Pub. Mus., Milwaukee, Wisconsin*.
- and —. 1988. West Indian amphibians and reptiles: a checklist. *Contrib. Biol. Geol. Milwaukee Pub. Mus.* (74):1-264.
- and —. 1991. Amphibians and reptiles of the West Indies: descriptions, distributions, and natural history. *Univ. Florida Press, Gainesville*.
- Shochat, D. and H.C. Dessauer. 1981. Comparative immunological study of albumins of *Anolis* lizards of the Caribbean islands. *Comp. Biochem. Physiol.* 68A:67-73.
- Telford, S.R., Jr. 1975. Saurian malaria in the Caribbean: *Plasmodium azurophilum* sp. nov., a malarial parasite with schizogony and gametogony in both red and white blood cells. *Intl. J. Parasitol.* 5:383-394.
- Underwood, G. and E. Williams. 1959. The anoline lizards of Jamaica. *Bull. Inst. Jamaica, Sci. Ser.*, (9):1-48.
- Williams, E.E. 1972. The origin of faunas. Evolution of lizard congeners in a complex island fauna: a trial analysis. *Evol. Biol.* 6:47-89.
- . 1976. West Indian anoles: a taxonomic and evolutionary summary. 1. Introduction and a species list. *Breviora* (440):1-21.
- . 1983. Ecomorphs, faunas, island size, and diverse end points in island radiations of *Anolis*, p. 326-370. *In* R.B. Huey, E.R. Pianka, and T.W. Schoener (eds.), *Lizard ecology: studies of a model organism*. Harvard Univ. Press, Cambridge, Massachusetts.
- and A.S. Rand. 1977. Species recognition, dewlap function and faunal size. *Amer. Zool.* 17:261-270.

Anthony P. Russell, Department of Biological Sciences, The University of Calgary, Calgary, Alberta, Canada T2N 1N4, and Aaron M. Bauer, Department of Biology, Villanova University, Villanova, Pennsylvania 19085, U.S.A.

Primary editor for this account, Andrew H. Price.

Published 15 October 1991 and Copyright © 1991 by the Society for the Study of Amphibians and Reptiles.